

Nexium 24HR Tablets

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredient (in each tablet)	Purpose
Esomeprazole 20 mg (Each delayed-release tablet corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer
Uses	
<ul style="list-style-type: none">■ treats frequent heartburn (occurs 2 or more days a week)■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
Warnings	
Allergy alert: Do not use if you are allergic to esomeprazole	
Do not use if you have:	
<ul style="list-style-type: none">■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools■ heartburn with lightheadedness, sweating or dizziness■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness■ frequent chest pain	
These may be signs of a serious condition. See your doctor.	
Ask a doctor before use if you have	
<ul style="list-style-type: none">■ had heartburn over 3 months. This may be a sign of a more serious condition.■ frequent wheezing, particularly with heartburn■ unexplained weight loss■ nausea or vomiting■ stomach pain	
Ask a doctor or pharmacist before use if you are taking	
<ul style="list-style-type: none">■ warfarin, clopidogrel or cilostazol (blood-thinning medicines)■ prescription antifungal or anti-yeast medicines■ digoxin (heart medicine)■ diazepam (anxiety medicine)■ tacrolimus or mycophenolate mofetil (immune system medicines)■ prescription antiretrovirals (medicines for HIV infection)■ methotrexate (arthritis medicine)	
Stop use and ask a doctor if	
<ul style="list-style-type: none">■ your heartburn continues or worsens■ you need to take this product for more than 14 days■ you need to take more than 1 course of treatment every 4 months■ you get diarrhea■ you develop a rash or joint pain	

Nexium 24HR Tablets

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- swallow whole. Do not crush or chew tablets.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age; ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, croscopovidone, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, mica, microcrystalline cellulose, paraffin, polyethylene glycol, polysorbate 80, sodium stearyl fumarate, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at **1-866-226-1600**